DRAFT VERSION FEBRUARY 14, 2013
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1302.2913v1 [astro-ph.HE] 12 Feb 2013

HELIUM-IGNITED VIOLENT MERGERS AS A UNIFIED MODEL FOR NORMAL AND RAPIDLY DECLINING TYPE IA SUPERNOVAE

R. PAKMOR[1,3], M. KROMER[2,3] AND S. TAUBENBERGER[2]

Draft version February 14, 2013

ABSTRACT

The progenitors of Type Ia Supernovae (SNe Ia) are still not known, despite significant progress during the last years in theory and observations. Violent mergers of two carbon–oxygen (CO) white dwarfs (WDs) are one candidate suggested to be responsible for at least a significant fraction of normal SNe Ia. Here, we propose a new, robust, explosion mechanism for violent mergers that works for CO primary WDs with CO as well as helium (He) WD companions. In this mechanism a detonation in the leftover He shell on top of the primary CO WD triggers a second detonation in its core, similar to the well-known double-detonation scenario for helium-accreting CO WDs. However, in contrast to that scenario where a massive He shell is required to form a spontaneous detonation, in the proposed scenario the He detonation is ignited dynamically by a merger of the binary system or instabilities in the accretion stream. Accordingly the required helium-shell mass is significantly smaller, and hence its burning products are unlikely to affect the optical display of the explosion. We show that this scenario has the potential to explain the different brightness distributions, delay times and relative rates of normal and fast declining SNe Ia. Finally, we discuss extensions to our unified merger model needed to obtain a comprehensive picture of the full observed diversity of SNe Ia.

Subject headings: Supernovae: general, white dwarfs, binaries: general

1. INTRODUCTION

Historically, Type Ia supernovae (SNe Ia) have been believed to be a uniform class of objects. However, over the past two decades large supernova surveys have revealed that this class is in fact rather inhomogeneous. Nonetheless, until recently, it has been argued that most – if not all – SNe Ia originate from delayed detonations of Chandrasekhar-mass carbon–oxygen (CO) White Dwarfs (WDs) accreting hydrogen-rich material from a non-degenerate companion star (e.g. Mazzali et al. 2007).

Lately, however, an increasing number of observational constraints have challenged this idea. In particular, the very nearby normal SN Ia 2011fe did not show any signs of a companion star (Li et al. 2011; Bloom et al. 2012), nor any signature of hydrogen in nebular spectra (Shappee et al. 2013) which is expected in the single-degenerate scenario owing to material stripped from the companion (Marietta et al. 2000; Leonard 2007). Moreover, there is a growing number of SN Ia remnants in which no remaining companion star could be found (e.g. Schaefer & Pagnotta 2012; González Hernández et al. 2012). Also, population-synthesis studies predict that the expected rate of single-degenerate Chandrasekhar-mass explosions is at least a factor of a few below the observed rate (e.g. Ruiter et al. 2009; Mennekens et al. 2010, but see Han & Podsiadlowski 2004 for a different result).

Double-degenerate systems, in contrast, avoid most of these problems: there is no surviving companion star expected and they can naturally explain the absence of hydrogen features in late-time spectra. At the same time, recent first-principle three-dimensional simulations of violent mergers of two CO WDs reproduce subluminous and normal SNe Ia well (Pakmor et al. 2010, 2012; Röpke et al. 2012).

Here, we discuss a new idea for the explosion mechanism in the violent-merger scenario and show how it enables us to construct a comprehensive picture for SNe Ia.

The paper is outlined as follows. In Sect. 2 we discuss the proposed explosion mechanism for violent mergers. We then describe how this naturally leads to two distinct populations of SNe Ia in Sect. 3. Sect. 4 presents the expected brightness distributions, rates and delay times of both populations in the context of observational constraints. In Sect. 5 we discuss some peculiar sub-classes of SNe Ia and how they can be integrated in our unified model for SNe Ia. We conclude with a brief summary in Sect. 6.

2. VIOLENT MERGERS REVISITED

In a violent merger of two CO WDs (Pakmor et al. 2010), a carbon detonation is assumed to form on the surface of the primary (more massive) WD when the secondary is about to be destroyed by tidal interactions and mass transfer heats up the surface of the primary to very high temperatures. The ensuing detonation burns and completely unbinds the merging object.

Although violent mergers are able to reproduce many observational characteristics of subluminous and normal SNe Ia (Pakmor et al. 2010, 2012; Röpke et al. 2012), it is debated whether a carbon detonation can really form under the temperature and density conditions found on the surface of the primary WD (Pakmor et al. 2010; Dan et al. 2011; Raskin et al. 2012; Pakmor et al. 2012; Seitenzahl et al. 2009). However, forming a helium (He) detonation requires much lower temperatures and densities compared to a carbon detonation (Seitenzahl et al. 2009), which are easily reached in the simulations.

Since most CO WDs have a small but non-negligible He shell (10^{-3} to $10^{-2}\,\mathrm{M}_{\odot}$, Iben & Tutukov 1985), the He will start to burn during the final stages of the merger, and a detonation may emerge. We argue that this He detonation forms either by the impact of the secondary on the primary WD

---

[1] Heidelberger Institut für Theoretische Studien, Schloss-Wolfsbrunnenweg 35, D-69118 Heidelberg, Germany

[2] Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany

[3] Kavli Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa, Chiba 277-8583, Japan



Figure 1. In the helium-ignited violent merger model a detonation forms in the thin He shell of a CO WD, either by the impact of the secondary WD or by accretion instabilities. Once a helium-shell detonation has been ignited, a converging shock runs through the CO core and triggers a secondary CO detonation that completely disrupts the system. The proposed mechanism works both for CO- and He-WD companions (as indicated by the two branches in the figure).

when the secondary is about to be destroyed (see e.g. Pakmor et al. 2012) or already before by instabilities in the accretion stream if mass transfer becomes dynamical (Guillochon et al. 2010). In both cases, the detonation forms shortly after mass transfer sets in, i.e. before a significant amount of mass can be transferred from the companion to the primary.

Once a helium-shell detonation has been ignited, a second detonation in the CO core is unavoidable when the shock from the shell detonation converges in the core (Fink et al. 2007). This mechanism has been shown to work down to helium-shell masses of only a few times $10^{-3}\,\mathrm{M}_{\odot}$ (Fink et al. 2010), thus providing a much more robust ignition mechanism for violent mergers than the previously discussed carbon surface ignition. Moreover, it works independently of the nature of the secondary (He or CO WD) and even for rather small mass ratios of the binary system (a sketch of the helium-ignited violent-merger model is shown in Fig. 1).

3. THE OBSERVATIONAL DISPLAY OF VIOLENT MERGERS

The explosion mechanism described above directly suggests that there are at least two different populations of SNe Ia, which originate from explosions in pairs of CO WDs and CO/He WDs, respectively. For the subsequent discussion it is irrelevant whether the explosion is caused by a merger or instabilities in the accretion stream, since they only differ slightly in the distance between primary and secondary WD at the time of the explosion.

In both populations, the amount of ^{56}Ni synthesized in the explosion, and therefore the intrinsic brightness, correlates directly with the mass of the CO primary WD, since the central density of the secondary WD is always below the threshold for ^{56}Ni production at the time of detonation. However, depending on the nature of the secondary WD, the total mass of the ejecta changes. In addition, details of the density profile and composition structure of the ejecta depend on the interaction of the the ashes of the primary WD with the secondary.

Carbon-ignited violent mergers of two CO WDs have already been shown to reproduce the light curves and spectra of normal SNe Ia fairly well (Pakmor et al. 2012; Röpke et al. 2012). Whether this holds for helium-ignited violent mergers has to be investigated by detailed simulations. In helium-accreting sub-Chandrasekhar-mass double-detonation models the spectral properties are highly sensitive to the burning products of the He shell (e.g. Kromer et al. 2010). However, in these models it is assumed that the He shell grows massive enough to become dynamically unstable and ignite spontaneously (Bildsten et al. 2007). In contrast, in our merger model, for a given primary WD mass the He shell is less massive than in those models. Therefore, it is conceivable that in our scenario the ashes of the He shell will have little or no effect on spectra and light curves.

It has also been shown that mergers of two low-mass CO WDs reproduce colors and spectra of objects similar to SN 1991bg very well (Pakmor et al. 2010). However, such mergers fail to reproduce the narrow, fast evolving light curves of 1991bg-like SNe. The light curve width of SNe Ia is mainly set by the amount of ^{56}Ni synthesized during the explosion since the ^{56}Ni also provides a significant part of the total opacity of the ejecta. As a consequence brighter explosions have typically more slowly declining light curves. For a given ^{56}Ni mass, however, the total ejecta mass will also affect the light curve width. We therefore propose that the progenitors of fast declining SNe Ia (most notably 1991bg-like objects) might be binaries of He and CO WDs. The narrow light curves could then be understood as a natural consequence of the smaller ejecta mass. At the same time, the spectral properties and colors of the explosion could be expected not to change drastically compared to a merger of two CO WDs with the same primary mass, since the secondary WD, which is burned after the primary, only affects the central parts of the ejecta (Pakmor et al. 2012).

4. BRIGHTNESS DISTRIBUTIONS, RATES AND DELAY TIMES

To explain the observed diversity of SNe Ia with a theoretical model, it is necessary to reproduce not only light curves and spectra of single objects, but also the brightness distribution of objects of different sub-classes, their relative rates, and their delay times.

In this regard, we base the analysis of our scenario on one specific population-synthesis model that has been described in a set of papers (Ruiter et al. 2009, 2011, 2013), since it is possibly the most detailed model published at the moment. Note that other population-synthesis models find partly to vastly different results, which introduces significant uncertainties and may alter some of our conclusions.

Ruiter et al. (2013) have shown that carbon-ignited violent



Figure 2. Observed diversity of SNe Ia in $\Delta m_{15}(B)$ vs. $M_{B,\mathrm{peak}}$ space as found in a sub-sample of the data presented in tables 8 and 9 of Hicken et al. (2009) where SNe without a $\Delta m_{15}(B)$ measurement and a distance modulus $\mu \leq 33$ mag have been removed. Most of these SNe Ia are normal and follow the Phillips et al. (1999) relation. Additional datapoints (Phillips et al. 2007; Taubenberger et al. 2008, 2011; Krisciunas et al. 2009; Maguire et al. 2011; Ganeshalingam et al. 2012) have been included to indicate the position of the peculiar sub-classes discussed in the text. Except for 2002cx-like SNe, our unified merger model potentially explains the full observed diversity. Explosions in CO/CO binaries are expected to have more-slowly declining light curves than those in CO/He binaries (indicated by the background-color gradient).

mergers of two CO WDs can not only explain the brightness distribution of normal SNe Ia but also their observed delay-time distribution and rates. Moreover, they found a tendency that mergers with more massive primary WDs, which lead to brighter supernovae, preferentially occur in younger populations, consistent with observed trends (see e.g. Gupta et al. 2011). However, Ruiter et al. (2013) find not enough systems to account for the observed number of faint SNe. In nature those have predominantly fast light curves similar to SN 1991bg (see Fig. 2), but there seems to emerge a sparse population with low luminosities yet slowly evolving light curves similar to SN 2002es and PTF10ops (Maguire et al. 2011; Ganeshalingam et al. 2012, see also Fig. 2) which might be consistent with the mergers of low-mass CO WD binaries.

In contrast to mergers of two CO WDs, the distribution of mass-transferring CO/He WD binaries is found to have a strong peak between 0.8 $M_\odot$ and 0.9 $M_\odot$ for the mass of the primary WD (Ruiter et al. 2011). Although stable mass transfer is assumed for the binary systems discussed in Ruiter et al. (2011), we argue that a significant fraction of them may either merge (Dan et al. 2011) or cause a He detonation due to instabilities in the accretion stream (Guillochon et al. 2010). Since this, in case, happens soon after mass transfer sets in, the core mass of a CO WD in Ruiter et al. (2011) is still a good indicator for the absolute brightness of the subsequent explosion. In general, binaries with a He WD secondary should therefore lead to fainter explosions since their primaries are typically significantly less massive than those of CO WD binaries.

Mergers with a primary WD mass around 0.9 $M_\odot$ should produce a ^{56}Ni mass very close to that required for 1991bg-like SNe Ia (Pakmor et al. 2010; Sim et al. 2010). For those primary masses, the rate of potentially exploding CO/He WD binaries dominates over that of pure CO WD binaries (Ruiter et al. 2011). Since these systems will also have narrow light curves (see discussion above) they would naturally explain the population of 1991bg-like SNe. Moreover, the very long delay times of these explosions (>1 Gyr; Ruiter et al. 2011) are in good agreement with the observed tendency of 1991bg-like SNe to arise in old stellar populations.



Figure 3. Histogram of the observed number of SNe Ia as a function of $\Delta m_{15}(B)$ (data from table 9 of (Hicken et al. 2009)). The distribution shows some indication for bimodality which might be identified with the two different companion populations in our model (illustrated by the background-color gradient as in Fig. 2).

Population synthesis calculations also find a sparsely populated tail of CO/He WD binaries towards larger primary masses (Ruiter et al. 2011). Explosions of such systems can reach luminosities more typical for normal SNe Ia but should still have faster light curves similar to 1991bg-like SNe. Observationally, these events might be identified with objects like SN 2003gs or SN 2007au (Krisciunas et al. 2009; Hicken et al. 2009, see Fig. 2).

It is interesting to note that the two populations may also provide a natural explanation for the observed bimodal distribution of SNe Ia in $\Delta m_{15}(B)$-space which some SN data sets suggest (see Fig. 3).

5. A COMPREHENSIVE PICTURE FOR TYPE IA SUPERNOVAE

Having presented a model that claims to explain a large part of the observed diversity of SNe Ia, we now investigate if other sub-classes identified in the literature can also be incorporated to obtain a comprehensive picture.

5.1. *Type Ia Supernovae with CSM interaction*

There is evidence that some fraction of SNe Ia show signs of CSM interaction. So far, this has mostly been interpreted as support for the single-degenerate scenario where recurrent nova outbursts are expected to produce shell-like density enhancements in the outflows from a companion star (Patat et al. 2011).

However, also the violent merger scenario predicts a range of observable CSM-interaction features, depending on the delay between the last common envelope (CE) and the supernova explosion. As discussed in Ruiter et al. (2013), about 0.5% of the violent mergers of two CO WDs explode less than

1000 years after the last CE event. Therefore, for these objects the material expelled in the CE phase is still close to the explosion site and interaction is unavoidable (see, e.g. Livio & Riess 2003).

If the delay between the last CE phase and the explosion is not too large there are three different regimes of CSM interaction in violent mergers. In most cases, the CE material will only be visible in absorption, potentially producing variable absorption-line features.[4] In rare systems with very short post-CE delays, the explosion ejecta will interact directly with the CE material, giving rise to hydrogen emission lines as observed for PTF11kx (Dilday et al. 2012). Finally, if the violent merger happens already during the CE phase, interaction will considerably contribute to the light curve and the explosion may appear as a Type IIn supernova at certain phases (cf. SN 2002ic; Livio & Riess 2003).

Since the contribution of the inspiral by gravitational-wave emission to the total delay time for violent mergers with CSM interaction is very small, we predict that they only occur in young stellar populations and should be absent in elliptical galaxies. In contrast, the traditional interpretation as single-degenerate symbiotic systems predicts long delay times (Hachisu et al. 2008).

5.2. *Superluminous Type Ia Supernovae*

Superluminous SNe Ia, which are significantly brighter than normal SNe Ia, have been a puzzle to explain so far (e.g. Howell et al. 2006). Recently, there is increasing support for the idea that they may be normal SNe Ia enshrouded with a CO envelope that contributes light at early times and produces dust at very late times (Hachinger et al. 2012; Taubenberger et al. submitted).

Such a structure might be explained by mergers of two massive CO WDs which do not have sufficient He available to form a detonation in the He shell of the primary WD, if this shell exists at all. If the carbon then does not ignite directly during the merger, the disrupted secondary will form an accretion disk and a hot extended envelope around the primary WD (Shen et al. 2012). Accretion onto the central WD might cause it to explode as a SN Ia a short time later (Zhu et al. 2012).

5.3. *2002cx-like Type Ia Supernovae*

In our picture, the vast majority of SNe Ia can be explained in terms of double-degenerate progenitors, without the need for explosions of Chandrasekhar-mass WDs. 2002cx-like SNe Ia, however, are different since they require strongly mixed ejecta (Jha et al. 2006) that cannot be produced in a detonation. As shown recently, pure deflagrations of Chandrasekhar-mass CO WDs ignited off-center reproduce 2002cx-like SNe Ia well (Kromer et al. 2013). Note that these explosions do not completely burn the WD, but leave behind a compact remnant (Kromer et al. 2013; Fink et al., in prep.).

6. SUMMARY

In this paper, we have introduced a new ignition scenario for double-degenerate systems in which a violent merger or accretion instability causes a thin He shell on the surface of the primary CO WD to detonate immediately after the onset of mass transfer. Like in the double-detonation explosion scenario for helium-accreting sub-Chandrasekhar-mass WDs, this He detonation subsequently leads to a detonation of the underlying CO core, which completely unbinds the progenitor stars.

The new scenario works independently of the nature of the secondary (He or CO) WD. We have argued that the two types of progenitor systems lead to two distinct populations of SNe Ia: mostly faint, fast declining objects for He companions and bright, slowly declining SNe for CO companions. This may naturally explain the observed bi-modality of SNe Ia in $\Delta m_{15}(B)$ space. We have also shown that population synthesis qualitatively predicts brightness distributions, relative rates and delay times for these two populations in agreement with observational data.

Finally, we have combined our new scenario with a recent model for 2002cx-like objects (Kromer et al. 2013) to construct a comprehensive picture of SNe Ia that is in principle able to explain all major sub-classes including superluminous SNe Ia and SNe Ia with CSM interaction.

R.P. and M.K. are grateful for IPMU's hospitality during a research visit at which this work was initiated. We thank A. Ruiter, M. Fink, S. Hachinger, W. Hillebrandt, F. Röpke, I. Seitenzahl and S. Sim for many inspiring discussions. This work was supported by the DFG through the cluster of excellence 'Origin and Structure of the Universe' (EXC 153) and the Transregional Collaborative Research Centre 'The Dark Universe' (TRR 33) and in part by World Premier International Research Center Initiative, MEXT, Japan.

REFERENCES

Bildsten, L., Shen, K. J., Weinberg, N. N., & Nelemans, G. 2007, ApJ, 662, L95
Bloom, J. S., Kasen, D., Shen, K. J., et al. 2012, ApJ, 744, L17
Dan, M., Rosswog, S., Guillochon, J., & Ramirez-Ruiz, E. 2011, ApJ, 737, 89
Dilday, B., Howell, D. A., Cenko, S. B., et al. 2012, Science, 337, 942
Fink, M., Hillebrandt, W., & Röpke, F. K. 2007, A&A, 476, 1133
Fink, M., Röpke, F. K., Hillebrandt, W., et al. 2010, A&A, 514, A53
Ganeshalingam, M., Li, W., Filippenko, A. V., et al. 2012, ApJ, 751, 142
González Hernández, J. I., Ruiz-Lapuente, P., Tabernero, H. M., et al. 2012, Nature, 489, 533
Guillochon, J., Dan, M., Ramirez-Ruiz, E., & Rosswog, S. 2010, ApJ, 709, L64
Gupta, R. R., D'Andrea, C. B., Sako, M., et al. 2011, ApJ, 740, 92
Hachinger, S., Mazzali, P. A., Taubenberger, S., et al. 2012, MNRAS, 427, 2057
Hachisu, I., Kato, M., & Nomoto, K. 2008, ApJ, 683, L127
Han, Z., & Podsiadlowski, P. 2004, MNRAS, 350, 1301
Hicken, M., Challis, P., Jha, S., et al. 2009, ApJ, 700, 331
Howell, D. A., Sullivan, M., Nugent, P. E., et al. 2006, Nature, 443, 308
Iben, Jr., I., & Tutukov, A. V. 1985, ApJS, 58, 661
Jha, S., Kirshner, R. P., Challis, P., et al. 2006, AJ, 131, 527
Krisciunas, K., Marion, G. H., Suntzeff, N. B., et al. 2009, AJ, 138, 1584
Kromer, M., Sim, S. A., Fink, M., et al. 2010, ApJ, 719, 1067
Kromer, M., Fink, M., Stanishev, V., et al. 2013, MNRAS, 429, 2287
Leonard, D. C. 2007, ApJ, 670, 1275
Li, W., Bloom, J. S., Podsiadlowski, P., et al. 2011, Nature, 480, 348
Livio, M., & Riess, A. G. 2003, ApJ, 594, L93
Maguire, K., Sullivan, M., Thomas, R. C., et al. 2011, MNRAS, 418, 747
Marietta, E., Burrows, A., & Fryxell, B. 2000, ApJS, 128, 615
Mazzali, P. A., Röpke, F. K., Benetti, S., & Hillebrandt, W. 2007, Science, 315, 825
Mennekens, N., Vanbeveren, D., De Greve, J. P., & De Donder, E. 2010, A&A, 515, A89
Miszalski, B., Jones, D., Rodríguez-Gil, P., et al. 2011, A&A, 531, A158

---

[4] Future observations of CE remnants are required to investigate whether such objects can reproduce the structure of distinct CSM shells as observed for e.g. SN 2006X (Patat et al. 2007). Potential candidates for CE remnants are planetary nebulae with central binary systems (e.g. Miszalski et al. 2011).

Pakmor, R., Kromer, M., Röpke, F. K., et al. 2010, Nature, 463, 61
Pakmor, R., Kromer, M., Taubenberger, S., et al. 2012, ApJ, 747, L10
Patat, F., Chugai, N. N., Podsiadlowski, P., et al. 2011, A&A, 530, A63
Patat, F., Chandra, P., Chevalier, R., et al. 2007, Science, 317, 924
Phillips, M. M., Lira, P., Suntzeff, N. B., et al. 1999, AJ, 118, 1766
Phillips, M. M., Li, W., Frieman, J. A., et al. 2007, PASP, 119, 360
Raskin, C., Scannapieco, E., Fryer, C., Rockefeller, G., & Timmes, F. X. 2012, ApJ, 746, 62
Röpke, F. K., Kromer, M., Seitenzahl, I. R., et al. 2012, ApJ, 750, L19
Ruiter, A. J., Belczynski, K., & Fryer, C. 2009, ApJ, 699, 2026
Ruiter, A. J., Belczynski, K., Sim, S. A., et al. 2011, MNRAS, 417, 408
Ruiter, A. J., Sim, S. A., Pakmor, R., et al. 2013, MNRAS, 429, 1425
Schaefer, B. E., & Pagnotta, A. 2012, Nature, 481, 164
Seitenzahl, I. R., Meakin, C. A., Townsley, D. M., Lamb, D. Q., & Truran, J. W. 2009, ApJ, 696, 515
Shappee, B. J., Stanek, K. Z., Pogge, R. W., & Garnavich, P. M. 2013, ApJ, 762, L5
Shen, K. J., Bildsten, L., Kasen, D., & Quataert, E. 2012, ApJ, 748, 35
Sim, S. A., Röpke, F. K., Hillebrandt, W., et al. 2010, ApJ, 714, L52
Taubenberger, S., Hachinger, S., Pignata, G., et al. 2008, MNRAS, 385, 75
Taubenberger, S., Benetti, S., Childress, M., et al. 2011, MNRAS, 412, 2735
Zhu, C., Chang, P., van Kerkwijk, M., & Wadsley, J. 2012, ArXiv e-prints